

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings of claims in the application:

Listing of Claims:

1-7. (Canceled)

8. (Currently amended) A method of visibly reducing a human skin wrinkle comprising: topically applying to the human skin wrinkle an imiquimod composition-IRM compound that is an agonist of TLR7, TLR8, or both TLR7 and TLR8 in an amount and for a period of time sufficient to visibly reduce the wrinkle, wherein imiquimod is the sole active pharmaceutical ingredient applied to the human skin wrinkle in performance of the method; wherein the IRM compound is an imidazoquinoline amine, a tetrahydroimidazoquinoline amine, an imidazopyridine amine, a 1,2 bridged imidazoquinoline amine, a 6, 7 fused cycloalkylimidazopyridine amine, an imidazonaphthyridine amine, a tetrahydroimidazonaphthyridine amine, an oxazoloquinoline amine, a thiazoloquinoline amine, an oxazolopyridine amine, a thiazolopyridine amine, an oxazonaphthyridine amine, a thiazolonaphthyridine amine, or a combination thereof.

9-32. (Canceled)

33. (New) The method of claim 8, wherein the imiquimod composition is applied daily.

34. (New) The method of claim 8, wherein the imiquimod composition is applied once per day two to three times a week.

35. (New) The method of claim 8, wherein the imiquimod composition comprises 5% imiquimod by weight.

1 36. (New) The method of claim 8, wherein the imiquimod composition about
2 1% to about 2.5% by weight.

1 37. (New) The method of claim 8, wherein the imiquimod composition is
2 applied to the human skin wrinkle for about one month.

1 38. (New) The method of claim 8, wherein the imiquimod composition is
2 applied to the human skin wrinkle for about two months.

1 39. (New) The method of claim 8, wherein the imiquimod composition is
2 applied to the human skin wrinkle for about 4 to 12 months.

1 40. (New) The method of claim 8, further comprising measuring the
2 reduction in human skin wrinkles by visual or photographic examination of the skin.

1 41. (New) The method of claim 8, wherein the imiquimod composition is
2 about 5% imiquimod by weight, and the imiquimod composition is applied to the human skin
3 wrinkle two to three times a week.

1 42. (New) The method of claim 8, wherein the imiquimod composition is
2 about 1% imiquimod by weight, and the imiquimod composition is applied daily to the human
3 skin wrinkle.

1 43. (New) The method of claim 8, wherein the imiquimod composition is
2 about 2.5% imiquimod by weight, and the imiquimod composition is applied daily to the human
3 skin wrinkle.